

Prevalence of Vitamin D Deficiency Among Healthy Infants and Toddlers

Catherine M. Gordon, MD, MSc; Henry A. Feldman, PhD; Linda Sinclair, BA; Avery LeBoff Williams, BA; Paul K. Kleinman, MD; Jeannette Perez-Rossello, MD; Joanne E. Cox, MD

Objectives: To determine the prevalence of vitamin D deficiency and to examine whether 25-hydroxyvitamin D (25OHD) concentration varies as a function of skin pigmentation, season, sun exposure, breastfeeding, and vitamin D supplementation.

Design: Cross-sectional sample.

Setting: Urban primary care clinic.

Participants: Healthy infants and toddlers (N=380) who were seen for a routine health visit.

Outcome Measures: Primary outcomes were serum 25OHD and parathyroid hormone levels; secondary measures included data on sun exposure, nutrition, skin pigmentation, and parental health habits. Wrist and knee radiographs were obtained for vitamin D-deficient participants.

Results: The prevalence of vitamin D deficiency (≤ 20 ng/mL) was 12.1% (44 of 365 participants), and 146 par-

ticipants (40.0%) had levels below an accepted optimal threshold (≤ 30 ng/mL). The prevalence did not vary between infants and toddlers or by skin pigmentation. There was an inverse correlation between serum 25OHD and parathyroid hormone levels (infants: $r = -0.27$, $P < .001$; toddlers: $r = -0.20$, $P = .02$). In multivariable models, breastfeeding without supplementation among infants and lower milk intake among toddlers were significant predictors of vitamin D deficiency. In vitamin D-deficient participants, 3 participants (7.5%) exhibited rachitic changes on radiographs, whereas 13 (32.5%) had evidence of demineralization.

Conclusions: Suboptimal vitamin D status is common among otherwise healthy young children. Predictors of vitamin D status vary in infants vs toddlers, information that is important to consider in the care of these young patients. One-third of vitamin D-deficient participants exhibited demineralization, highlighting the deleterious skeletal effects of this condition.

Arch Pediatr Adolesc Med. 2008;162(6):505-512

THERE HAS BEEN CONCERN raised about a resurgence of vitamin D deficiency and rickets among infants and children, with reports emerging in the United States from Alaska,^{1,2} Iowa,³ Nevada,⁴ California,⁵ North Carolina,⁶ Texas,⁷ and mother-infant pairs in Boston,⁸ among others.⁹ The prevalence of vitamin D deficiency in young children also appears to be high in other countries, including England,¹⁰ Greece,¹¹ and Canada.^{12,13} One study from China found a 65.3% prevalence of vitamin D deficiency among 12- to 24-month-olds, but few cases (3.7%) of radiographic or clinical rickets were noted.¹⁴ Previous studies suggest risk factors to be dark skin pigmentation^{1,3-12} and breastfeeding without supplementation.^{1-7,9,12,13} To date, reports have focused primarily on young infants compared with toddlers.

Thus, there appears to be less information available regarding variables that predispose young children to vitamin D deficiency as they wean to fortified milk and solid foods.

*For editorial comment
see page 583*

In 2004, another study reported a high prevalence of vitamin D deficiency among otherwise healthy teenagers who were seen for primary care.¹⁵ In the current study, we sought to determine the prevalence of vitamin D deficiency and radiographic rickets among young children seen at an urban primary care clinic. The primary objective was to test the hypothesis that vitamin D deficiency (25-hydroxyvitamin D [25OHD] level, ≤ 20 ng/mL [to convert to nanomoles per liter, multiply by

Author Affiliations: Divisions of Adolescent Medicine (Dr Gordon and Mss Sinclair and Williams) and Endocrinology (Drs Gordon and Feldman), Clinical Research Program (Dr Feldman), Department of Radiology (Drs Kleinman and Perez-Rossello), and Division of General Pediatrics (Dr Cox), Children's Hospital Boston, Boston, Massachusetts.

2.496]) is prevalent among healthy infants and toddlers to the same degree as among adolescents (ie, 42%).¹⁵ The secondary objective was to test the hypothesis that 25OHD concentrations are lower and PTH levels are higher during winter, especially among those with darker skin pigmentation, as was seen among adolescents.¹⁵ We sought to identify nutritional and lifestyle variables that represented predictors of vitamin D deficiency and 25OHD level. We also tested the hypothesis that breastfeeding without supplementation among infants, consumption of juice rather than milk in toddlers, dark skin pigmentation, and winter season would emerge as significant predictors of vitamin D deficiency. Last, among those identified to be vitamin D deficient biochemically, we documented the prevalence of rickets.

METHODS

STUDY PARTICIPANTS

We studied 380 primary care patients (age range, 8-24 months) who were seen for physical examinations between October 17, 2005, and June 15, 2007, at the Children's Hospital Boston Primary Care Center and were undergoing a routine blood draw. The center is hospital based and provides well-child and acute care for 12 000 children annually, approximately 80% of whom live within the adjacent urban community. The population is primarily African American and Latino, and approximately 25% of parents are immigrants from diverse regions. The poverty rate for the communities served is 22% to 27%, compared with 20% for the city of Boston.¹⁶ Exclusion criteria included presence of a chronic disease and use of medications known to affect vitamin D metabolism during the previous 3 months. Parents or guardians provided written informed consent at study enrollment. The Committee on Clinical Investigation, Children's Hospital Boston, approved the protocol.

CLINICAL AND ANTHROPOMETRIC DATA

Data were collected at the clinic visit, including age, sex, height, weight, nutritional intake, daily sun exposure, and pertinent findings (eg, genu varum) from the physical examination. Skin color and sensitivity were evaluated by a research assistant (L.S. or A.L.W.) using established methods.^{17,18} The skin pigmentation scale ranged from 1 (heavily pigmented, eg, black) to 4 (lightly pigmented, eg, white). The skin sensitivity scale ranged from 1 (burns easily) to 6 (never burns).

NUTRITIONAL AND LIFESTYLE QUESTIONNAIRES

Each parent or guardian completed a questionnaire regarding nutritional intake for the parent and child. For children 1 year and older, the questionnaire covered consumption of milk, juice, fortified cereal, and water. Parents of infants younger than 1 year provided a breastfeeding history. The nutritional questionnaire asked parents to recall intake of vitamin D-fortified cereal, seafood, and milk. Parents also were asked about their own diets, use of vitamin D or other supplements, time spent outdoors, socioeconomic status, race/ethnicity, and education level.

LABORATORY MEASUREMENTS

One blood sample (15 mL) was obtained for each participant. Blood tests were performed in the hospital laboratory or sent

to ARUP Laboratories (Salt Lake City, Utah). Serum 25OHD levels were determined at ARUP Laboratories using a chemiluminescent assay (LIAISON; DiaSorin Inc, Stillwater, Minnesota). Serum calcium, phosphorus, magnesium, and alkaline phosphatase levels were measured locally using an end point assay in a multichannel analyzer (Roche Diagnostics, Indianapolis, Indiana). Intact PTH was measured by a 2-site chemiluminescence immunoassay (Nichols Institute, San Clemente, California). Samples were analyzed in multiple assays. Interassay coefficients of variation were 5.4% to 7.0% for PTH, 8.6% to 10.0% for 25OHD, 0.7% for alkaline phosphatase, and 1.5% to 2.2% for the cations.

DEFINITIONS

The patients were divided into 3 diagnostic categories according to serum 25OHD levels, as rounded to the nearest integer: vitamin D deficiency (≤ 20 ng/mL), severe vitamin D deficiency (≤ 8 ng/mL), and suboptimal vitamin D status (≤ 30 ng/mL). The definition of vitamin D deficiency was based on growing consensus among experts that a 25OHD level of 20 ng/mL or less is the appropriate diagnostic threshold for deficiency.¹⁹⁻²¹ This threshold also represents the low end of the reference range provided by the manufacturer of the assay. There is agreement among many skeletal health experts that 30 ng/mL represents an optimal level for 25OHD²²⁻²⁴ or a threshold of adequate concentration. This is supported by the fact that serum 25OHD level has been shown to be inversely correlated with PTH level at concentrations of 30 to 40 ng/mL.^{20,25,26} We decided a priori that a 5% prevalence of vitamin D deficiency would be clinically significant. The definition of severe vitamin D deficiency was based on the sensitivity of the 25OHD assay (7 ng/mL).

RADIOGRAPHIC EVALUATION

Participants found to have vitamin D deficiency underwent frontal bilateral wrist and knee computed radiography. Soft copies of radiographs were reviewed for rachitic changes on a single high-resolution monitor by 2 pediatric radiologists (P.K.K. and J.P.-R.), each working from a unique randomized list of participants. The interrater correlation coefficient was calculated for the Thacher et al²⁷ and demineralization scales. Each radiologist interpreted all images independently. Evidence of rickets on wrist and knee radiographs (eg, metaphyseal fraying) was determined using a validated 10-point scoring system.²⁷ The score progressed in half-point increments from 0 (normal) to 10 (severe) and included both wrists and knees. Scores from the 2 radiologists were averaged, and data on interrater reliability was recorded. A scale (0-5) was also used by each radiologist to assess degree of demineralization, and the scores were averaged.

STATISTICAL ANALYSIS

We used the Fisher exact test to compare the prevalence of vitamin D deficiency between infants and toddlers and to identify binary or polytomous variables associated with vitamin D deficiency within each patient group. To identify continuous variables associated with vitamin D deficiency, we used simple logistic regression with deficiency as the dichotomous dependent variable. To identify binary or polytomous variables associated with 25OHD level, we used the *t* test or 1-way analysis of variance (Fisher F test). To identify continuous variables associated with 25OHD level, we used simple linear regression.

We constructed a multiple logistic regression model for vitamin D deficiency and a multiple linear regression model for 25OHD concentration using all predictors of interest, whether or not the simple association was significant. We tested for con-

founding and masking relationships by adding or removing variables and observing the effect on statistical significance of the remaining variables in the model. The results were corroborated by forward and backward stepwise procedures for automatic variable selection.

We used Pearson *r* to determine the correlation between 25OHD and PTH levels and to characterize other relationships among variables. We used the intraclass correlation coefficient [(patient variance)/(patient + rater + residual variance)] to determine interrater reliability for radiographic scales. Statistical analyses were conducted with SAS statistical software, version 9.1 (SAS Institute Inc, Cary, North Carolina).

RESULTS

CLINICAL CHARACTERISTICS

The final sample included 380 subjects (**Table 1**). Biochemical measurements were available for 365 (**Table 2**). Among toddlers, mean intake of milk exceeded that of juice (Table 1). Cation concentrations were within the reference range for 354 infants and toddlers; 1 participant (0.2%) had hypocalcemia, 5 (1.4%) had hypercalcemia, 1 (0.2%) had hypophosphatemia, and 4 (1.1%) had hypermagnesemia. Elevated PTH levels were seen in 5 (1.4%).

DESCRIPTIVE CHARACTERISTICS OF CAREGIVERS

Of 380 infants, 377 (99.2%) were accompanied to the clinic visit by their mother (mean [SD] age, 28 [7] years). Forty-six adults (12.1%) were white, 233 (61.3%) were African American, 110 (28.9%) were Latino, 18 (4.7%) were Asian, and 10 (2.6%) were of another race or ethnicity. A total of 254 (66.8%) reported receiving governmental assistance. Ninety-three (24.5%) reported multivitamin use and 262 (68.9%) reported at least occasional consumption of fortified milk. In addition, 290 (76.3%) reported spending at least a half hour outdoors daily, but only 68 (17.9%) reported sunscreen use.

PREVALENCE OF VITAMIN D DEFICIENCY

The prevalence of vitamin D deficiency (≤ 20 ng/mL) was 44 of 365 children (12.1%) for the total sample, with 7 (1.9%) having severe deficiency (≤ 8 ng/mL); 146 (40.0%) had levels below an accepted optimal threshold (≤ 30 ng/mL). The prevalence did not differ between infants and toddlers (Table 1; $P > .30$ for all criteria). Participants identified as vitamin D deficient were invited to participate in a treatment trial. Those who chose not to participate were referred to their primary care provider for treatment.

VARIABLES ASSOCIATED WITH DEFICIENCY OR 25OHD LEVEL

For infants and toddlers, there was a modest, but significant, inverse correlation between serum 25OHD and PTH levels (infants: $r = -0.27$, $P < .001$; toddlers: $r = -0.20$, $P = .02$). There were no significant differences in hormone levels or the strength of the correlation between the variables in infants vs toddlers.

Table 1. Characteristics of Participants^a

Characteristic	Infants (n=247)	Toddlers (n=133)
Sex		
Girls	131 (53.0)	61 (45.9)
Boys	116 (47.0)	72 (54.1)
Season of visit		
Spring	58 (23.5)	29 (21.8)
Summer	60 (24.3)	34 (25.6)
Autumn	55 (22.3)	41 (30.8)
Winter	74 (30.0)	29 (21.8)
Skin pigmentation score		
1, Heavily pigmented	141 (57.1)	75 (56.4)
2	57 (23.1)	29 (21.8)
3	31 (12.6)	11 (8.3)
4, Lightly pigmented	18 (7.3)	18 (13.5)
Uses sunscreen	51 (21.6)	42 (33.3)
Skin sensitivity score		
1, Burns easily	4 (1.6)	1 (0.8)
2, Burns always	7 (2.8)	7 (5.3)
3, Burns moderately	32 (13.0)	10 (7.5)
4, Burns minimally	53 (21.5)	25 (18.8)
5, Burns rarely	99 (40.1)	56 (42.1)
6, Never burns	52 (21.1)	34 (25.6)
Feeding		
Breast, no vitamin D supplement	14 (5.7)	NA
Breast, vitamin D supplement	6 (2.4)	NA
Bottle and breast	42 (17.0)	NA
Bottle only	185 (74.9)	NA
Serum 25OHD level, ng/mL ^b		
≤ 8	4 (1.7)	3 (2.4)
≤ 20	26 (10.8)	18 (14.4)
≤ 30	97 (40.4)	49 (39.2)
Mean (SD) [range]		
Age, mo	9.6 (0.8) [7.6-12.7]	15.7 (3.6) [12.0-24.0]
Weight, kg	9.1 (1.2) [5.4-13.0]	10.8 (1.6) [7.5-14.6]
Body mass index ^c	17.7 (1.5) [13.6-22.3]	17.4 (1.9) [12.9-24.8]
Time spent outdoors, h/d	1.3 (1.1) [0-3]	1.4 (1.1) [0-3]
Milk intake, cups per day	NA	2.6 (1.8) [0-9]
Juice intake, cups per day	NA	2.0 (1.8) [0-10]
Fortified cereal intake, cups per day	NA	0.7 (0.7) [0-3]

Abbreviations: NA, not applicable; 25OHD, 25-hydroxyvitamin D.

SI conversion factor: To convert 25OHD to nanomoles per liter, multiply by 2.496.

^aData are given as the number (percentage) of participants unless otherwise indicated.

^bCategories are nested; each dichotomy was tested separately.

^cCalculated as weight in kilograms divided by height in meters squared.

Among hypothesized predictors of 25OHD level, we found a marked effect for breastfeeding without supplementation in infants (**Figure**). In toddlers, 25OHD level increased 2.9 ng/mL per cup per day of milk consumption (95% confidence interval [CI], 1.4-4.4; $P < .001$) (Figure). In infants, sunny season was paradoxically associated with a lower 25OHD level (mean [SE], 31.1 [1.1] ng/mL from April to September vs 37.8 [1.1] ng/mL from October to March; $P < .001$).

Table 2. Laboratory Findings

Level	Mean (SD) [Range]		Infant Mean (SD)–Toddler Mean (SD)	P Value ^b
	Infants ^a	Toddlers ^a		
25OHD, ng/mL	34.3 (12.3) [7-72]	35.2 (15.2) [7-96]	-0.8 (1.5)	.59
PTH, pg/mL	30.3 (37.0) [5-508]	29.6 (30.9) [4-334]	0.7 (3.8)	.85
Alkaline phosphatase, U/L	313 (270) [106-2987]	352 (446) [100-4293]	-39 (38)	.37
Calcium, mg/dL	10.51 (0.43) [7.0-11.9]	10.39 (0.36) [9.6-11.3]	0.12 (0.04)	.005
Magnesium, mEq/L	1.95 (0.16) [1.5-2.5]	1.88 (0.16) [1.6-2.2]	0.15 (0.05)	<.001
Phosphorus, mg/dL	5.87 (0.51) [3.7-7.0]	5.69 (0.70) [2.5-7.2]	0.18 (0.06)	.01

Abbreviations: 25OHD, 25-hydroxyvitamin D; PTH, parathyroid hormone.

SI conversion factors: To convert alkaline phosphatase to microkatal per liter, multiply by 0.0167; calcium to millimoles per liter, multiply by 0.25; magnesium to millimoles per liter, multiply by 0.50; 25OHD to nanomoles per liter, multiply by 2.496; PTH to nanograms per liter, multiply by 0.1053; and phosphorus to millimoles per liter, multiply by 0.323.

^aSample size varied between 238 and 242 for infants and 125 and 127 for toddlers.

^bP value for equal distribution in infants and toddlers by *t* test (not assuming equal variance); confirmed by Wilcoxon rank sum test.




Figure. Nutritional factors associated with serum levels of 25-hydroxyvitamin D (25OHD) in 247 infants at their 9-month visit to a primary care clinic (A) and 133 toddlers at their 18-month visit (B). Bars indicate mean and standard error of the mean. D Suppl indicates vitamin D supplementation. To convert 25OHD to nanomoles per liter, multiply by 2.496.

Variables potentially associated with vitamin D deficiency were examined one at a time by Fisher exact test or simple logistic regression (**Table 3**). The prevalence of deficiency did not vary by sex, season, time spent outdoors, sunscreen use, sun sensitivity, or skin pigmentation. Breastfeeding without supplementation was strongly associated with vitamin D deficiency among infants, with a more than 10-fold increase in risk relative to infants who were exclusively bottle-fed (Table 3). Among toddlers, milk consumption significantly decreased odds of deficiency (odds ratio [OR], 0.51; 95% CI, 0.34-0.76; $P=.001$), whereas juice consumption had the opposite effect (OR, 1.31; 95% CI, 1.02-1.69; $P=.03$). There was no significant simple association between body mass index (calculated as weight in kilograms divided by height in meters squared) and vitamin D deficiency in either infants (OR, 0.84 per kg/m^2 ; 95% CI, 0.62-1.13; $P=.25$) or toddlers (OR, 1.05 per kg/m^2 ; 95% CI, 0.77-1.42; $P=.76$). In the toddler group, there was no significant association between fortified cereal intake and vitamin D deficiency (OR, 1.27 per cup per day; 95% CI, 0.62-2.60; $P=.51$).

INDEPENDENT PREDICTORS OF DEFICIENCY OR 25OHD LEVEL

Multiple logistic regression models, evaluating predictors of vitamin D deficiency with adjustment for other predictors, are summarized in **Table 4**. Among infants, breastfeeding without supplementation markedly increased the odds of vitamin D deficiency compared with infants who were exclusively bottle-fed. Among toddlers, milk consumption significantly decreased odds of deficiency, whereas body mass index showed an opposite, marginally significant effect. In the adjusted analysis, juice consumption showed no effect on vitamin D deficiency in toddlers.

These results were corroborated by multiple linear regression models for 25OHD level. Breastfeeding without supplementation in infants was associated with a significantly lower level of 25OHD (mean [SD], 16.7 [3.4] ng/mL vs 36.3 [0.9] ng/mL in infants who were exclusively bottle-fed; $P<.001$). In toddlers, milk consumption (3.1-ng/mL increase in 25OHD level per cup of milk

Table 3. Variables Potentially Associated with Vitamin D Deficiency

Predictor	Infants			Toddlers		
	No. of Participants	Vitamin D Deficiency, No. (%) ^a	P Value ^b	No. of Participants	Vitamin D Deficiency, No. (%) ^a	P Value ^b
All	240	26 (10.8)	NA	125	18 (14)	NA
Sex						
Girls	127	16 (12.6)	.41	55	8 (14.5)	.99
Boys	113	10 (8.8)		70	10 (14.3)	
Season of visit						
Spring	57	6 (10.5)	.99	27	4 (14.8)	.65
Summer	59	7 (11.9)		33	6 (18.2)	
Autumn	52	5 (9.6)		37	6 (16.2)	
Winter	72	8 (11.1)		28	2 (7.1)	
Skin pigmentation score						
1, Heavily pigmented	136	16 (11.8)	.82	67	9 (13.4)	.41
2	57	5 (8.8)		29	6 (20.7)	
3	29	4 (13.8)		11	0 (0)	
4, Lightly pigmented	18	1 (5.6)		18	3 (16.7)	
Taken outdoors						
Y	208	23 (11.1)	.99	111	17 (15.3)	.69
N	32	3 (9.4)		14	1 (7.1)	
Uses sunscreen						
Y	49	5 (10.2)	.99	40	6 (15.0)	.99
N	181	19 (10.5)		78	12 (15.4)	
Skin sensitivity score						
1, Burns easily	4	0 (0)	.46	1	1 (100.0)	.16
2, Burns always	6	0 (0)		7	0 (0)	
3, Burns moderately	32	4 (12.5)		10	1 (10.0)	
4, Burns minimally	52	4 (7.7)		25	5 (20.0)	
5, Burns rarely	95	15 (15.8)		51	9 (17.6)	
6, Never burns	51	3 (5.9)		31	2 (6.5)	
Feeding						
Breast, no vitamin D supplement	14	9 (64.3)	<.001	NA	NA	NA
Breast, vitamin D supplement	6	2 (33.3)		NA	NA	NA
Bottle and breast	42	4 (9.5)		NA	NA	NA
Bottle only	178	11 (6.2)		NA	NA	NA

Abbreviations: NA, not applicable; 25OHD, 25-hydroxyvitamin D.

^aVitamin D deficiency defined as 25OHD level ≤ 20 ng/mL (to convert to nanomoles per liter, multiply by 2.496).

^bP values for equal prevalence in all categories determined by Fisher exact test.

per day; 95% CI, 1.5-4.8; $P < .001$) and body mass index (-2.7 -ng/mL decrease in 25OHD level per kilogram per meters squared; 95% confidence interval, -4.5 to -0.8 ; $P = .005$) were significantly associated with 25OHD level. In infants, sunny season was paradoxically associated with lower 25OHD levels (mean [SD], 31.6 [1.1] ng/mL from April to September vs 37.8 [1.1] ng/mL from October to March; $P < .001$).

RADIOGRAPHIC ASSESSMENTS

Of the 44 participants who had vitamin D deficiency, 40 (90.9%) returned to undergo bilateral wrist and knee radiography. Using the Thacher system,²⁷ 3 (7.5%; 0.8% of the total sample) exhibited rachitic changes. One child had visible genu varum on physical examination and a total knee-wrist score of 2. Comparing results from the 2 radiologists, the interrater correlation coefficient for the total knee-wrist score was 0.69. Using the demineralization scale, 13 of 40 (32.5%; 3.5% of the total sample) had evidence of demineralization (range of scores, 1-3 on a 5-point scale). All participants with

rachitic changes also exhibited demineralization. The interrater correlation coefficient for this second assessment scale was 0.42.

COMMENT

We found a relatively high prevalence of vitamin D deficiency among otherwise healthy infants and toddlers in a convenience sample from an urban clinic. A striking 40% had serum 25OHD levels below a threshold that is becoming increasingly accepted as ideal for bone health and protection conferred against malignancy, infection, and other forms of disease.²²⁻²⁴ Among children found to be vitamin D deficient, a few exhibited rachitic changes radiographically, but one-third had evidence of demineralization. Only 1 child had signs of rickets on physical examination. Thus, these infants and toddlers had a subclinical deficiency that could make detection of this issue particularly problematic in routine clinical practice, as a child's vitamin D status is not typically evaluated as part of routine care. Among infants, breastfeeding with-

Table 4. Multivariable Models for Vitamin D Deficiency

Predictor	Infants		Toddlers	
	OR (95% CI) ^a	P Value ^b	OR (95% CI) ^a	P Value ^b
Boys vs girls	1.34 (0.46-3.95)	.59	1.31 (0.35-4.90)	.69
Sunny season (April-September)	2.12 (0.71-6.33)	.18	0.64 (0.16-2.49)	.51
Skin pigmentation, per scale point	0.69 (0.32-1.49)	.34	0.69 (0.32-1.47)	.33
Time spent outdoors, per hour	1.01 (0.61-1.66)	.98	0.87 (0.43-1.78)	.70
Sunscreen use	2.48 (0.74-8.32)	.14	0.74 (0.16-3.45)	.70
Skin sensitivity, per scale point	1.17 (0.63-2.17)	.62	0.55 (0.27-1.15)	.11
Feeding, vs bottle only				
Breast, no vitamin D supplement	74.8 (13.5-416)	<.001	NA	NA
Breast, vitamin D supplement	4.42 (0.43-45.4)	.21	NA	NA
Bottle and breast	1.32 (0.34-5.16)	.68	NA	NA
Body mass index ^c	1.05 (0.74-1.48)	.79	1.57 (0.97-2.54)	.07
Milk intake, per cup per day	NA	NA	0.37 (0.21-0.65)	<.001
Juice intake, per cup per day	NA	NA	1.14 (0.73-1.78)	.56
Fortified cereal intake, per cup per day	NA	NA	1.73 (0.72-4.15)	.22

Abbreviations: CI, confidence interval; NA, not applicable; 25OHD, 25-hydroxyvitamin D; OR, odds ratio.

^aOdds of vitamin D deficiency, defined as 25OHD level ≤ 20 ng/mL (to convert to nanomoles per liter, multiply by 2.496), compared with reference category; for continuous predictors, increase in odds per unit predictor. Estimate and 95% confidence interval from multiple logistic regression analysis, adjusted for all other predictors listed.

^bP value odds ratio equal to unity (no effect).

^cCalculated as weight in kilograms divided by height in meters squared.

out supplementation was an independent predictor of vitamin D deficiency, replicating data from previous studies.^{1-4,6,9,12,13} Among toddlers, a protective effect of milk consumption was seen in univariable and multivariable models. Unexpectedly, we found that skin pigmentation, sun exposure, and sunscreen use were not predictors of 25OHD concentration or vitamin D deficiency, as was hypothesized.

Vitamin D concentration was not found to vary by skin pigmentation or sun sensitivity score. These data differ from our study of adolescents¹⁵ and other studies of infants or young children, in which dark skin pigmentation was associated with vitamin D deficiency.^{1,3,4,6-12} Interestingly, our sample was enriched with 233 African American children, but skin pigmentation was not identified as a risk factor for deficiency. Instead, this problem was prevalent across all variants of skin pigmentation. Because infants are often swaddled in blankets or dressed in more layers of clothing than older children or teenagers, cutaneous vitamin D synthesis may have been decreased or prevented in our participants, even during sunnier months, although this explanation is speculative and was not formally assessed by this study. In addition, it is noteworthy that average temperatures are lower in northern states, which may lead to more coverage with ultraviolet light-blocking clothing and reduced cutaneous synthesis of vitamin D. Concerns about safety could have also contributed to less sun exposure in this urban sample. Skin sensitivity, skin pigmentation, time spent outdoors, and sunscreen use were examined, but none was found to be a predictor of vitamin D status.

Our finding of a high prevalence of vitamin D deficiency among breastfed infants is consistent with earlier reports.^{1-6,9,12,13} Interestingly, strong significant trends were seen despite the fact that exclusive breastfeeding was strikingly low among study participants. The current recom-

mendation of the American Academy of Pediatrics^{28,29} and the Institute of Medicine³⁰ is to provide 200 IU of vitamin D daily to most infants, children, and adolescents, although some have questioned whether this dosage recommendation is adequate.³¹⁻³³ In this study, only 2% of breastfed infants were receiving vitamin D supplementation. All Massachusetts Medicaid-based insurance plans cover multivitamins containing 200 IU of vitamin D, although supplemental vitamins are not available through the Special Supplemental Nutrition Program for Women, Infants, and Children. Thus, lack of supplementation may have been related to barriers to access or lack of awareness. Furthermore, the fact that some infants were vitamin D deficient despite reported supplementation raises questions about the current dosage recommendation and overall compliance. These issues will need to be addressed in future studies that include a larger subset of breastfed infants and more rigorous measures of compliance. Nonetheless, these findings endorse recommendations for health care providers and parents to ensure that breastfed infants receive daily vitamin D supplementation for the duration of breastfeeding.

The current study is strengthened by the fact that radiographic assessments, another biomarker of vitamin D status, were obtained in a subset of patients, in addition to biochemical measurements. Previous studies have commonly been case series of infants with known rickets.^{4,7,12,14} Demineralization was detected radiographically in one third of our vitamin D-deficient participants. Although most deficiencies were subclinical, radiographic data suggest that many children exhibited deleterious effects. One group has advocated that physicians obtain a 25OHD measurement and wrist radiographs as a screening measure for high-risk children,³⁴ with consideration given to the possibility of multiple risk factors being present in a given child. Examples include a breastfed infant with a low

serum 25OHD level who lives at a high latitude or an overweight African American toddler from the same northern city. Another approach would be to obtain radiographs for those with evidence of longstanding vitamin D deficiency (eg, low 25OHD level with secondary hyperparathyroidism and/or physical manifestations of rickets). The child in this study with both physical and radiographic evidence of rickets exhibited this biochemical profile, including an undetectable serum 25OHD level and a markedly elevated PTH level. Because rickets can cause significant morbidity, radiographic evaluation, in addition to biochemical screening, would be an appropriate consideration in the most at-risk child.

Among the toddlers studied, there was a significant inverse relationship noted between body mass index and 25OHD level in multivariable models. This association was noted in our previous study of adolescents¹⁵ and in previous studies of adults.³⁵⁻³⁷ In simple logistic regression models, only juice and milk consumption were significant predictors of vitamin D deficiency, suggesting that confounding existed between body mass index and juice consumption. One study has suggested that vitamin D deficiency in adults who are obese is caused by decreased bioavailability, because of deposition of this fat-soluble vitamin in adipose tissue.³⁷ The current data suggest that an elevated body mass index is a risk factor for vitamin D deficiency. Although there is no consensus regarding the appropriateness of routine vitamin D screening for overweight patients, daily vitamin D supplementation, as for all children, would be warranted. Higher supplementation doses may be needed in this circumstance, but longitudinal studies examining the relationship between specific doses of vitamin D supplementation and both vitamin D levels and other health outcomes are needed to answer that question.

These findings must be considered in light of acknowledged limitations. The study was cross-sectional; therefore, causality cannot be inferred. The study sample was enriched with subgroups known to be at higher risk for vitamin D deficiency, including African American and Latino children. Interestingly, however, we did not see the expected association of vitamin D deficiency with dark skin pigmentation. The strikingly low prevalence of breastfed infants in the sample also limits the generalizability of these results. Nonetheless, we observed a strong, significant association between breastfeeding and vitamin D deficiency, especially in the absence of vitamin D supplementation, that deserves attention. The fact that we used an urban convenience sample may also limit the generalizability of these findings. We obtained information on weekly outdoor activities, but this measure may have provided only indirect information about sun exposure, and we found no association between this variable and 25OHD level. We also did not ask parents about clothing and swaddling practices, information that would have been especially informative because we noted an unexpected significant trend of low 25OHD levels during sunny months. Although we asked whether parents and their children were receiving vitamin D supplementation, more specific information on the actual quantity of vitamin D received and compliance would assist in determining whether noncompliance affected our results. Finally, in-

formation on nutrition and health habits for children and caretakers was obtained by self-report, which has inherent limitations.

In summary, these data add to accumulating evidence that vitamin D deficiency, a preventable health problem, is common among infants and toddlers, across ethnicity and season. A relationship was noted between the risk of vitamin D deficiency and lack of supplementation among breastfed infants and among toddlers with a higher body mass index. Among toddlers, there was a protective effect seen between milk consumption and lower risk of deficiency. These data underscore the fact that all breastfed infants should receive vitamin D supplementation for the duration of breastfeeding. However, further studies are needed to determine whether a 200 IU daily dose will provide adequate supplementation. Given the potential benefits of vitamin D on bone³⁸ and other tissues,^{22-24,26} and growing data supporting its immunomodulatory and antiproliferative effects,³⁹⁻⁴³ the current findings support recommendations advocating for vitamin D supplementation for all young children.^{28-30,44} These findings also raise the important question of whether some children, such as those with established risk factors for vitamin D deficiency, should receive periodic measurements of 25OHD levels.

Accepted for Publication: October 28, 2007.

Correspondence: Catherine M. Gordon, MD, MSc, Divisions of Adolescent Medicine and Endocrinology, Children's Hospital Boston, 300 Longwood Ave, Boston, MA 02115 (catherine.gordon@childrens.harvard.edu).

Author Contributions: Dr Gordon had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. *Study concept and design:* Gordon, Kleinman, and Cox. *Acquisition of data:* Gordon, Sinclair, Williams, Kleinman, Perez-Rossello, and Cox. *Analysis and interpretation of data:* Gordon, Feldman, Sinclair, Kleinman, and Perez-Rossello. *Drafting of the manuscript:* Gordon, Feldman, Williams, and Kleinman. *Critical revision of the manuscript for important intellectual content:* Gordon, Feldman, Sinclair, Williams, Kleinman, Perez-Rossello, and Cox. *Statistical analysis:* Feldman. *Obtained funding:* Gordon. *Administrative, technical, or material support:* Sinclair, Williams, and Cox. *Study supervision:* Gordon and Cox.

Financial Disclosure: None

Funding/Support: This study was supported by grants from the Allen Foundation Inc and the McCarthy Family Foundation; grant NIH MO1-RR-2172 from the National Center for Research Resources (to the Children's Hospital Boston General Clinical Research Center); and project T71 MC00009 of the Maternal and Child Health Bureau, US Health Resources and Services Administration.

Role of the Sponsor: The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

Additional Contributions: Stephanie Bristol, BS, Melissa Chittle, BA, and Lisa Bartels, BA, provided excellent technical assistance. We also thank our patients and their families for their participation.

1. Gessner BD, deSchweinitz E, Petersen KM, Lewandowski C. Nutritional rickets among breast-fed black and Alaska Native children. *Alaska Med*. 1997;39(3):72-74.
2. Gessner BD, Plotnick J, Muth PT. 25-hydroxyvitamin D levels among healthy children in Alaska. *J Pediatr*. 2003;143(4):434-437.
3. Ziegler EE, Hollis BW, Nelson SE, Jeter JM. Vitamin D deficiency in breastfed infants in Iowa. *Pediatrics*. 2006;118(2):603-610.
4. Shaikh U, Alpert PT. Nutritional rickets in Las Vegas, Nevada. *J Pediatr Endocrinol Metab*. 2006;19(3):209-212.
5. Hayward I, Stein MT, Gibson MI. Nutritional rickets in San Diego. *Am J Dis Child*. 1987;141(10):1060-1062.
6. Kreiter SR, Schwartz RP, Kirkman HN, Charlton PA, Calikoglu AS, Davenport ML. Nutritional rickets in African American breast-fed infants. *J Pediatr*. 2000;137(2):153-157.
7. Shah M, Salhab N, Patterson D, Seikaly MG. Nutritional rickets still afflict children in north Texas. *Tex Med*. 2000;96(6):64-68.
8. Lee JM, Smith JR, Philipp BL, Chen TC, Mathieu J, Holick MF. Vitamin D deficiency in a healthy group of mothers and newborn infants. *Clin Pediatr (Phila)*. 2007;46(1):42-44.
9. Weisberg P, Scanlon KS, Li R, Cogswell ME. Nutritional rickets among children in the United States: review of cases reported between 1986 and 2003. *Am J Clin Nutr*. 2004;80(6)(suppl):1697S-1705S.
10. Callaghan AL, Moy RJ, Booth IW, Debelle G, Shaw NJ. Incidence of symptomatic vitamin D deficiency. *Arch Dis Child*. 2006;91(7):606-607.
11. Nicolaidou P, Hatzistamatiou Z, Papadopoulou A, et al. Low vitamin D status in mother-newborn pairs in Greece. *Calcif Tissue Int*. 2006;78(6):337-342.
12. Binet A, Kooh SW. Persistence of vitamin D-deficiency rickets in Toronto in the 1990s. *Can J Public Health*. 1996;87(4):227-230.
13. Ward LM, Gaboury I, Ladhani M, Zlotkin S. Vitamin D-deficiency rickets among children in Canada. *CMAJ*. 2007;177(2):161-166.
14. Strand MA, Perry J, Jin M, et al. Diagnosis of rickets and reassessment of prevalence among rural children in northern China. *Pediatr Int*. 2007;49(2):202-209.
15. Gordon CM, DePeter K, Feldman HA, Grace E, Emans SJ. Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adol Med*. 2004;158:531-537.
16. Boston Foundation. The Boston Indicators Project: Economy. <http://www.bostonindicators.org/IndicatorsProject/Economy/Indicator.aspx?id=2538>. Accessed October 30, 2007.
17. Pathak MA, Jimbow K, Szabo G, Fitzpatrick TB. Sunlight and melanin pigmentation. In: Goldsmith K, ed. *Photochemical and Photobiological Reviews*. New York, NY: Plenum Press; 1976:211-239.
18. Jimbow K, Fitzpatrick B, Wick MM. Biochemistry and physiology of melanin pigmentation. In: Goldsmith LA, ed. *Physiology, Biochemistry, and Molecular Biology of the Skin*. New York, NY: Oxford University Press; 1991:873-909.
19. Malabanan A, Veronikis IE, Holick MF. Redefining vitamin D insufficiency. *Lancet*. 1998;351(9105):805-806.
20. Thomas MK, Lloyd-Jones DM, Thadhani RI, et al. Hypovitaminosis D in medical inpatients. *N Engl J Med*. 1998;338(12):777-783.
21. Holick MF. High prevalence of vitamin D inadequacy and implications for health. *Mayo Clin Proc*. 2006;81(3):353-373.
22. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentration of 25-hydroxyvitamin D for multiple health outcomes [published correction appears in *Am J Clin Nutr*. 2006;84(5):1253]. *Am J Clin Nutr*. 2006;84(1):18-28.
23. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357(3):266-281.
24. Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. *Osteoporos Int*. 2005;16(7):713-716.
25. Chapuy MC, Preziosi P, Maamer M, et al. Prevalence of vitamin D insufficiency in an adult normal population. *Osteoporos Int*. 1997;7(5):439-443.
26. Holick MF, Siris ES, Binkley N, et al. Prevalence of vitamin D inadequacy among postmenopausal North American women receiving osteoporosis therapy. *J Clin Endocrinol Metab*. 2005;90(6):3215-3224.
27. Thacher TD, Fischer PR, Pettifor JM, Lawson JO, Manaster BJ, Reading JC. Radiographic scoring method for the assessment of the severity of nutritional rickets. *J Trop Med*. 2000;46(3):132-139.
28. Gartner LM, Greer FR; Section on Breastfeeding and Committee on Nutrition, American Academy of Pediatrics. Prevention of rickets and vitamin D deficiency: new guidelines for vitamin D intake. *Pediatrics*. 2003;111(4, pt 1):908-910.
29. Gartner LM, Morton J, Lawrence RA, et al; American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 2005;115(2):496-506.
30. Food and Nutrition Board. *Institute of Medicine: Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride*. Washington, DC: National Academy Press; 1997.
31. Heaney RP. Long-latency deficiency disease: insights from calcium and vitamin D. *Am J Clin Nutr*. 2003;78(5):912-919.
32. Heaney RP. Functional indices of vitamin D status and ramifications of vitamin D deficiency. *Am J Clin Nutr*. 2004;80(6)(suppl):1706S-1709S.
33. Weaver CM, Fleet JC. Vitamin D requirements: current and future. *Am J Clin Nutr*. 2004;80(6)(suppl):1735S-1739S.
34. Spence JT, Serwint JR. Secondary prevention of vitamin D-deficiency rickets. *Pediatrics*. 2004;113(1, pt 1):e70-e72. <http://pediatrics.aappublications.org/cgi/content/full/113/1/e70>.
35. Vilarrasa N, Maravall J, Estepa A, et al. Low 25-hydroxyvitamin D concentrations in obese women: their clinical significance and relationship with anthropometric and body composition variables. *J Endocrinol Invest*. 2007;30(8):653-658.
36. Bischof MG, Heinze G, Vierhapper H. Vitamin D status and its relation to age and body mass index. *Horm Res*. 2006;66(5):211-215.
37. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr*. 2000;72(3):690-693.
38. Zamora SA, Rizzoli R, Belli DC, Slosman DO, Bonjour JP. Vitamin D supplementation during infancy is associated with higher bone mineral mass in prepubertal girls. *J Clin Endocrinol Metab*. 1999;84(12):4541-4544.
39. EURODIAB Study Group. Vitamin D supplement in early childhood and risk for type 1 (insulin-dependent) diabetes mellitus. *Diabetologia*. 1999;42(1):51-54.
40. Hyppönen E, Laara E, Reunanen A, Jarvelin MR, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet*. 2001;358(9292):1500-1503.
41. Munger KL, Levin LI, Hollis BW, Howard NS, Ascherio A. Serum 25-hydroxyvitamin D levels and risk for multiple sclerosis. *JAMA*. 2006;296(23):2832-2838.
42. Holt PR, Arber N, Halmos B, et al. Colonic epithelial cell proliferation decreases with increasing levels of serum 25-hydroxyvitamin D. *Cancer Epidemiol Biomarkers Prev*. 2002;11(1):113-119.
43. Grant WB, Garland CF. A critical review of studies on vitamin D in relation to colorectal cancer. *Nutr Cancer*. 2004;48(2):115-123.
44. Calvo MS, Whiting SJ, Barton CN. Vitamin D fortification in the United States and Canada: current status and data needs. *Am J Clin Nutr*. 2004;80(6)(suppl):1710S-1716S.