

# Multi-institutional Profile of Adults Admitted to Pediatric Intensive Care Units

Jeffrey D. Edwards, MD, MA, MAS; Amy J. Houtrow, MD, MPH; Eduard E. Vasilevskis, MD; R. Adams Dudley, MD, MBA; Megumi J. Okumura, MD, MAS

**Importance:** Growing numbers of persons with childhood-onset chronic illnesses are surviving to adulthood. Many use pediatric hospitals for their inpatient needs. To our knowledge, the prevalence and characteristics of adult pediatric intensive care unit patients have not been reported.

**Objectives:** To estimate the proportion of adults admitted to pediatric intensive care units (PICUs), characterize them, and compare them with older adolescents.

**Design:** One-year cross-sectional analysis.

**Setting:** Pediatric intensive care units in the United States that participated in the Virtual Pediatric Intensive Care Unit Systems.

**Participants:** Pediatric intensive care unit patients 15 years or older admitted in 2008.

**Main Outcome Measures:** We compared adults with adolescents across clinical characteristics and outcomes. Mixed-effects logistic regression was used to estimate the independent association of age with PICU mortality.

**Results:** Seventy PICUs had 67 629 admissions; 1954 admissions (2.7%) were patients 19 years or older; and 9105 admissions (13.5%) were patients aged 15 to 18 years. The proportion of adults ( $\geq 19$  years) varied considerably by PICU (range, 0%-9.2%). As age increased, the proportion of patients who had a complex chronic condition and planned or perioperative admissions increased; the proportion of trauma-related admissions decreased. Patients aged 21 to 29 years had a 2 times (95% CI, 1.3-3.2;  $P = .004$ ) greater odds of PICU mortality compared with adolescent patients, after adjusting for Paediatric Index of Mortality score, sex, trauma, and having a complex chronic condition. Being 30 years or older was associated with a 3.5 (95% CI, 1.3-9.7;  $P = .01$ ) greater odds of mortality.

**Conclusions and Relevance:** In this multi-institutional study, adults constituted a small but high-risk proportion of patients in some PICUs, suggesting that these PICUs should have plans and protocols specifically focused on this group.

*JAMA Pediatr.* 2013;167(5):436-443.

Published online March 18, 2013.

doi:10.1001/jamapediatrics.2013.1316

**G**ROWING NUMBERS OF PERSONS with childhood-onset chronic illnesses are surviving to adulthood. Pediatric and adult medicine stakeholders have highlighted the importance of transitioning young adults with special health care needs from child- to adult-oriented health care.<sup>1-4</sup> Transitioning care may occur at different ages, depending on many factors, including access to appropriate adult care and patient preference. Although a focus on outpatient care is essential for successful transition,<sup>4,5</sup> the transition of inpatient care is also necessary. However, some patients with childhood-onset chronic illnesses, such as congenital heart disease, cystic fibrosis, and sickle cell disease, continue to receive inpatient care from pediatric hospitals well into adulthood.<sup>6-11</sup>

While adults presenting to pediatric emergency departments or admitted to pediatric hospitals have been described,<sup>6,12</sup> to our knowledge, the extent to which and

*For editorial comment  
see page 485*

under what circumstances adults are admitted to pediatric intensive care units (PICUs) has not been addressed. Neither has there been a description of their clinical characteristics or outcomes. This information will improve our understanding of how this population impacts PICU case mix, outcome performance measures, and organizational needs and help clinicians to better tailor strategies for transitioning patients to adult services. To characterize adults admitted to US PICUs and compare them with adolescents, we

Author Affiliations are listed at the end of this article.

conducted a retrospective multi-institutional cohort study of PICU patients 15 years and older.

## METHODS

### DATA SOURCE

We performed an analysis of all patients 15 years and older who were admitted in 2008 to 70 US PICUs that participated in the Virtual Pediatric Intensive Care Unit Systems (VPS). The VPS is a collaboration between the Children's Hospital Association, National Outcomes Center of the Children's Hospital and Health System in Wisconsin, and Children's Hospital Los Angeles. The VPS database contains encounter-level information on all admissions entered by trained clinicians at the participating sites. Sites submit data for their medical/surgical PICUs and pediatric cardiac intensive care units (ICUs); data are not collected from neonatal ICUs, intermediate or specialized non-critical care units, or general wards. Annual certification of data definitions, routine interrater reliability testing, and automated and manual data cleaning queries are used to ensure data quality. All institutions are required to submit several demographic and clinical data elements. Some sites voluntarily submitted additional characteristics. These included race (46 sites), secondary diagnoses (68 sites), primary reason for admission (49 sites), trauma (55 sites), airway/ventilatory procedures (32 sites), and functionality scores (21 sites). Data for mandatory characteristics were submitted for every admission by each site. Similarly, when a site elected to report a nonmandatory characteristic, data were submitted for every admission.

On approval of our research proposal, data were provided by VPS. No endorsement or editorial restriction of the interpretation of these data or opinions of the authors have been implied or stated. We received an exemption from the University of California, San Francisco Committee on Human Research for this study.

### PATIENT CHARACTERIZATION

Patients were stratified into 4 age categories: 15 to 18, 19 to 20, 21 to 29, and 30 years or older. These age categories were chosen because we considered 15 to 18 years the upper end of childhood. Nineteen to 20 years can be considered a transition period, while 21- to 29-year-old patients are adults sometimes still cared for by pediatric providers. We considered 30 years or older beyond young adulthood and the age range for the pediatric health care system. We presented the aggregate number and proportion of PICU patients and admissions, as well as the median proportions with interquartile ranges (IQRs) among the VPS sites.

We collected patient characteristics of each PICU admission. When information was available for only a subgroup of the admissions and sites, we noted this in the text or Tables. Demographic characteristics included age, sex, and race. Clinical characteristics included type of admission (unplanned, perioperative, or trauma), reason for admission by primary organ system affected, patient origin and disposition, whether admission was to a PICU with more than 1000 annual admissions, whether positive pressure ventilation was used, severity of illness, and presence of any complex chronic condition. Positive pressure ventilation was defined as conventional mechanical ventilation, high-frequency ventilation, and bilevel or continuous positive airway pressure via tracheal, mask, or nasal interface. Severity of illness was assessed using a predicted probability of mortality estimated with the Paediatric Index of Mortality 2.<sup>13</sup> Complex chronic conditions were defined according to the Feudtner et al definition<sup>14</sup> but were identified among VPS diagnoses using a list of VPS

codes we developed and detailed elsewhere.<sup>15</sup> Using data from PICUs that provided secondary diagnoses, we identified specific high-morbidity childhood-onset chronic conditions that are no longer lethal in childhood but require continuing management in adulthood. Using data from PICUs that provided procedural information, we also identified procedures and operations classified by organ system.

We also assessed outcomes of care: functional status, length of stay, and mortality. Functional outcomes were assessed with preadmission and discharge Pediatric Overall and Cerebral Performance Categories (POPC and PCPC)<sup>16</sup> from those units that reported this information. The POPC and PCPC scores range from 1 (normal function) to 6 (brain death). Scores of 2, 3, and 4 indicate mild, moderate, and severe disability, respectively. A score of 5 indicates coma or vegetative state. Length of stay was measured in days from time of PICU admission to PICU discharge. Finally, we assessed PICU mortality.

### STATISTICAL ANALYSES

We calculated proportions and 95% confidence intervals or medians and IQR for demographic and clinical characteristics, stratified by age categories. Similarly, we calculated proportions and 95% confidence intervals for specific chronic and acute conditions or categories, as well as for procedures and operations classified by organ system. Bivariate statistics were calculated using Pearson  $\chi^2$  test or Kruskal-Wallis rank test when appropriate.

We estimated the association between age and PICU mortality by calculating a prevalence risk ratio and the risk difference for PICU mortality of patients 19 years or older compared with those aged 15 to 18 years. To estimate the adjusted association, we then fitted a mixed-effects logistic regression model, using the adolescent and adult cohort from units that supplied secondary diagnoses and trauma data. Random effects were at the hospital level. Our primary predictors were the age categories, and we adjusted for sex, trauma, severity of illness (Paediatric Index of Mortality 2) score, and presence of any complex chronic condition. Model diagnostics were performed to ensure accuracy and goodness of fit. Statistical significance was determined using a *P* value of  $<.05$  and by constructing 95% confidence intervals. Stata version 12 (StataCorp) was used for statistical analyses.

## RESULTS

### SITE AND PATIENT CHARACTERISTICS

In 2008, 70 VPS PICUs were located in 32 states; these PICUs represented approximately 30% of the Children's Hospital Association–designated children's hospitals in the United States.<sup>17</sup> Fifteen PICUs (21%) had fewer than 10 beds, 20 (44%) had 11 to 20 beds, and 25 (35%) had 21 or more beds. Twenty-eight PICUs (40%) had more than 1000 annual admissions. Twenty-nine PICUs (41%) were situated in freestanding children's hospitals; 58 (83%) were level I units.<sup>18</sup> Fifty-three PICUs (76%) were affiliated with a pediatric residency program and 28 (40%), with a critical care fellowship.

These PICUs had 50 786 patients with 67 629 admissions in 2008. Of these, 1954 admissions (2.7%) were patients 19 years or older; 9105 admissions (13.5%) were patients aged 15 to 18 years. Further breakdown of number of admissions by age group is shown in **Table 1**. The oldest patient was 56 years of age. The proportion of adults admitted to each PICU varied widely; the median pro-

**Table 1. Patient Characteristics and Functional Status by Age Groups**

Characteristic	% of Admissions (95% CI)			
	Age 15-18 y (n = 9105)	Age 19-20 y (n = 1102)	Age 21-29 y (n = 6520)	Age ≥30 y (n = 100)
Admissions	83.1 (82.4-83.8)	10.1 (9.5-10.6)	5.9 (5.5-6.4)	0.9 (0.7-1.1)
Male	54.4 (53.3-55.4)	51.5 (48.6-54.5)	55.2 (51.4-59)	43 (33.1-52.9)
Race <sup>a</sup>				
White	58.7 (57.5-59.9)	61.1 (57.8-64.4)	78.3 (74.5-82.2)	82.4 (73.1-91.6)
African American	16.7 (15.9-17.8)	16.8 (14.3-19.4)	8.9 (6.3-11.6)	2.9 (0-7.1)
Hispanic	15.4 (14.6-16.3)	13.5 (11.2-15.9)	8.9 (6.3-11.6)	7.4 (1-13.7)
Asian/Indian/Pacific Islander	2.7 (2.3-3.1)	3.4 (2.2-4.6)	1.1 (0.4-2.1)	
Other/unspecified	6.3 (5.7-6.9)	5.1 (3.6-6.5)	2.7 (1.2-4.2)	7.4 (1-13.7)
Had a complex chronic condition <sup>b</sup>	52.7 (51.6-53.7)	76.5 (74-79.1)	83.8 (80.9-86.7)	79.3 (70.9-87.8)
PIM2 risk of mortality, median % (IQR)	0.8 (0.3-1.4)	0.9 (0.4-2.2)	1 (0.4-3.2)	1 (0.3-2)
Functional status, median (IQR) <sup>c</sup>				
POPC score, preadmission	1 (1-2)	2 (1-3)	3 (1-4)	2 (1-3)
POPC score, discharge	2 (1-3)	2 (1-3)	3 (1-4)	2 (1-3)
PCPC score, preadmission	1 (1-1)	1 (1-2)	2 (1-3)	1 (1-3)
PCPC score, discharge	1 (1-1)	1 (1-3)	2 (1-4)	1 (1-3)

Abbreviations: IQR, interquartile range; PCPC, Pediatric Cerebral Performance Category; PIM2, Paediatric Index of Mortality 2; POPC, Pediatric Overall Performance Category.

<sup>a</sup>Based on 7923 admissions (72%) of patients older than 15 years from 46 units that supplied race.

<sup>b</sup>Based on 10 481 admissions (96%) of patients older than 15 years from 68 units that supplied secondary diagnoses.

<sup>c</sup>Based on 3542 admissions (32%) of patients older than 15 years from 21 units that supplied POPC and PCPC scores.

portion of admissions for patients 19 years or older was 1.8% (IQR, 0.5%-3.2%; range, 0%-9.2%); aged 19 to 20 years, 1% (IQR, 0.4%-2.1%); and 21 years or older, 0.6% (IQR, 0%-1%).

Additional patient characteristics by age category are presented in Table 1. Admission characteristics and PICU outcomes are presented in **Table 2**. As the age category of admissions increased, there were higher proportions of patients who were white, had a complex chronic condition, and had planned or perioperative admissions. Also, as age category increased, there were lower proportions of patients who were admitted for trauma-related reasons.

## CONDITIONS AND PROCEDURES

The proportions of specific chronic and acute conditions among our study admissions are presented in **Table 3** and **Table 4**, respectively. Several chronic conditions were more common among adults than adolescent PICU patients, including congenital cardiac abnormalities (13.5% vs 4.8%;  $P < .001$ ), generalized developmental delay (10.4% vs 4.7%;  $P < .001$ ), and epilepsy (12.9% vs 7.4%;  $P < .001$ ). Acute lung injury and respiratory distress syndrome were more common in adults (18.1%) compared with adolescents (11.4%;  $P < .001$ ). The proportions of admissions for patients who underwent procedures and operations by age category are presented in **Table 5**.

## OUTCOMES

The POPC and PCPC scores were available from 21 units for 3542 admissions (32%) and are presented in Table 1. In general, adults had worse preadmission functionality, as evidenced by higher median overall scores (moderate disability vs no disability in adolescents) or wider distributions of POPC and PCPC scores. On discharge,

the median POPC score of adolescents was worse (mild disability) than preadmission POPC score (no disability). The preadmission and discharge POPC scores did not change for adults.

The PICU mortality for the entire cohort was 2.7%; mortality increased significantly with age categories ( $P < .001$ ). Adults 19 years or older had a 126% higher incidence of PICU mortality than adolescent patients did (risk ratio, 2.3; 95% CI, 1.8-2.9; risk difference, 2.8%; 95% CI, 1.8%-3.9%). To address the impact of trauma and having a complex chronic condition on mortality, we used a subgroup of 8828 admissions (81%) from 63 units (overall mortality, 2.6%) that supplied secondary diagnoses and trauma data to fit a regression model of PICU mortality. After adjusting for these variables plus Paediatric Index of Mortality 2 score and sex, patients aged 21 to 29 years had a 2.02 times (95% CI, 1.26-3.24;  $P = .004$ ) greater odds of PICU mortality compared with adolescents. Being 30 years or older was associated with a 3.55 (95% CI, 1.30-9.73;  $P = .01$ ) greater adjusted odds of PICU mortality. Age 19 to 20 years was not statistically associated with PICU mortality (adjusted odds ratio, 1.34;  $P = .21$ ) compared with age 15 to 18 years. Finally, as age category increased, the median length of stay also increased significantly ( $P < .001$ ). The median length of stay for adults (1.9 days) was half a day longer than for adolescents (1.4 days).

## COMMENT

To our knowledge, this is the first multi-institutional study to characterize adults admitted to PICUs. Commonly, adult patients are excluded from studies that describe pediatric inpatient populations and from most pediatric inpatient databases. We found that approximately 2% of PICU patients and 3% of admissions from this cohort were

**Table 2. Admission Characteristics and PICU Outcomes by Age Groups**

Characteristic	% of Admissions (95% CI)			
	Age 15-18 y (n = 9105)	Age 19-20 y (n = 1102)	Age 21-29 y (n = 652)	Age ≥30 y (n = 100)
Origin				
OR/PACU/procedure suite	33.2 (32.2-34.2)	38.7 (35.9-41.6)	43.3 (39.4-47.1)	60 (50.2-69.8)
Emergency department	48.8 (47.7-49.8)	32.6 (29.8-35.3)	30.2 (26.7-33.7)	14 (7.1-20.9)
General ward	11.9 (11.2-12.6)	20.1 (17.7-22.4)	16.4 (13.6-19.3)	14 (7.1-20.9)
Another ICU	1.8 (1.5-2.1)	2.7 (1.8-3.7)	2.6 (2.4-3.8)	2 (0-4.8)
Intermediate unit	1.5 (1.2-1.7)	1.6 (0.9-2.4)	1.8 (0.8-2.9)	5 (0.7-9.3)
Chronic/rehabilitation facility	0.3 (0.2-0.4)	0.4 (0-0.7)	1.4 (0.5-2.3)	2 (0-4.8)
Outpatient/home	2.3 (2-2.6)	3.8 (2.7-4.9)	4.1 (2.6-5.7)	2 (0-4.8)
Other	0.3 (0.1-0.4)	0.1 (0-0.3)	0.2 (0-0.5)	1 (0-3)
Reason for admission <sup>a</sup>				
Respiratory	21.2 (20.3-22.2)	30.6 (27.5-33.6)	36.8 (32.6-40.9)	25.6 (20-32.3)
Neurologic	25.3 (24.3-26.3)	20.8 (18.1-23.5)	17.8 (14.5-21.1)	7.3 (1.6-13.1)
Cardiovascular/shock	14.4 (13.6-15.2)	19.1 (16.5-21.7)	22.6 (19-26.2)	32.9 (22.5-43.3)
Metabolic	13.4 (12.6-14.1)	8 (6.2-9.8)	4.8 (3-6.7)	3.7 (0-7.8)
Hemorrhage/coagulopathy	3.5 (3.1-3.9)	3.2 (2.1-4.4)	2.7 (1.3-4.1)	6.1 (0.8-11.4)
Procedure	1 (0.8-1.2)	2.5 (1.5-3.6)	1.9 (0.7-3.1)	...
Other	21.3 (20.3-22.2)	15.7 (13.3-18.2)	13.3 (10.4-16.3)	24.4 (14.9-33.9)
Admitted to PICU with >1000 annual admissions	64.1 (63.1-65)	72.1 (69.5-74.8)	80.4 (77.3-83.4)	90 (84-96)
Unplanned admission	72.6 (71.7-73.5)	67.8 (65-70.5)	63.3 (59.6-67.1)	44 (34.1-53.9)
Preoperative/postoperative	37.1 (36.1-38.1)	41.7 (38.7-44.6)	47.7 (43.8-51.5)	60 (50.2-69.8)
Trauma <sup>b</sup>	11.4 (10.6-12.1)	3.5 (2.2-4.7)	2.1 (0.8-3.4)	1.5 (0-4.6)
Used PPV <sup>c</sup>	25 (23.9-26.2)	33 (29.4-36.7)	45.8 (40.6-51)	49 (34.8-63.2)
Disposition				
General ward	66.7 (65.8-67.7)	65.7 (62.9-68.5)	54.1 (50.3-58)	38 (28.3-47.7)
Another ICU	1.2 (1-1.4)	1.8 (1-2.6)	2.1 (1-3.3)	3 (0-6.4)
Intermediate unit	8.6 (8.1-9.2)	9.5 (7.8-11.2)	14 (11.3-16.6)	32 (22.7-41.3)
OR	0.4 (0.3-0.5)	0.4 (0-0.7)	0.2 (0-0.4)	...
Chronic/rehabilitation facility	0.9 (0.7-1.1)	1.2 (0.5-1.8)	2.9 (1.6-4.2)	2 (0-4.8)
Home	16.7 (15.9-17.4)	16.3 (14.1-18.5)	19.6 (16.6-22.7)	18 (10.3-25.7)
Died	2.2 (1.9-2.5)	4 (2.8-5.2)	6.7 (4.8-8.7)	6 (1.3-10.7)
Other	3.2 (2.8-3.6)	1.1 (0.6-1.9)	0.3 (0-1.1)	1 (0-5.4)
PICU LOS, d, median (IQR)	1.4 (0.9-2.9)	1.8 (1-3.9)	2 (1-4.9)	2.1 (1-4.3)
PICU mortality	2.2 (1.9-2.5)	4 (2.8-5.2)	6.7 (4.8-8.7)	6 (1.3-10.7)

Abbreviations: ellipses, no admissions; ICU, intensive care unit; IQR, interquartile range; LOS, length of stay; OR, operating room; PACU, postanesthesia care unit; PICU, pediatric intensive care unit; PIM, Pediatric Index of Mortality; PPV, positive pressure ventilation.

<sup>a</sup>Based on 8525 admissions (78%) of patients older than 15 years from 49 units that supplied primary reason for admission.

<sup>b</sup>Based on 8831 admissions (81%) of patients older than 15 years from 55 units that supplied the trauma variable.

<sup>c</sup>Based on 6443 admissions (59%) of patients older than 15 years from 32 units that supplied information on respiratory support.

adults 19 years or older. There was wide variation in the proportion of adult patients across PICUs, from zero to 9%. Adults were generally admitted to PICUs with higher patient volume in greater proportions. Adult PICU patients had greater severity of illness and preadmission disability, longer lengths of stay, and higher mortality compared with younger patients. We also observed that the majority of adult admissions were planned and perioperative, and the proportion appeared to be higher with older corresponding age.

The proportion of adult patients in our PICU cohort is similar to the general hospital population presented by Goodman et al.<sup>10</sup> They showed that 1.8% of discharges from the inpatient services of 41 tertiary children's hospitals were patients older than 18 years and 0.8% of patients and 1% of discharges were older than 21 years. These adults also had higher rates of chronic conditions compared with younger patients. Our study adds to the current literature by focusing on PICU patients, characterizing their type of and reasons for PICU admissions, and describing adult patients' chronic conditions in greater detail.

After age 18 years, close to 80% of PICU patients had a complex chronic condition, compared with 53% of adolescents. These chronic conditions increase a person's risk for critical illness and need for critical care.<sup>15</sup> Notably, many (20%) adult PICU patients 21 years or older had congenital cardiac abnormalities and/or had cardiac procedures or operations. Goodman et al<sup>10</sup> reported an even larger proportion of adult inpatients with congenital heart disease (29%). Cerebral palsy, generalized developmental delay, and epilepsy were other relatively common chronic conditions in our adult cohort. The higher proportion of adults with certain childhood-onset chronic conditions may reflect the inability or unwillingness to transition to adult care or greater need for critical care services as certain conditions progress. Lower proportions may indicate more successful transitioning of certain populations or higher mortality at older ages in others (eg, muscular dystrophy).

Although this study cannot determine the reasons for persistent care of adults in PICUs, the disproportionate number with chronic conditions suggests some adults with

**Table 3. Common Chronic Conditions by Age Groups**

Condition	% of Admissions (95% CI) <sup>a</sup>			
	Age 15-18 y (n = 8741)	Age 19-20 y (n = 1036)	Age 21-29 y (n = 612)	Age ≥30 y (n = 92)
Cardiac abnormality, congenital	4.9 (4.4-5.3)	9.6 (7.8-11.3)	15.6 (12.8-18.6)	46.7 (36.3-57.1)
Chromosomal anomaly/genetic abnormality	2.3 (1.9-2.6)	5.2 (3.9-6.6)	6 (4.2-7.9)	4.3 (0.1-8.6)
Gastrointestinal				
Chronic liver disease	0.7 (0.5-0.8)	1.9 (1.1-2.7)	1 (0.2-1.8)	4.3 (0.1-8.6)
Gastrostomy	2.6 (2.3-3)	5.7 (4.3-7.1)	10.8 (8.3-13.2)	8.7 (2.8-14.6)
Other childhood-onset conditions	0.1 (0-0.2)	0.4 (0-0.8)	0.5 (0-1)	...
Hematologic				
Sickle cell anemia	1 (0.8-1.2)	2.1 (1.2-3)	1.1 (0.3-2)	...
Other childhood-onset conditions	0.3 (0.2-0.4)	1.3 (0.6-1.9)	...	...
Immunologic				
Human immunodeficiency virus infection	0.2 (0.1-0.2)	0.4 (0-0.8)	...	...
Other childhood-onset conditions	0.1 (0.1-0.2)	0.8 (0.2-1.3)	0.2 (0-0.5)	...
Malignancy				
Bone marrow transplant	1 (0.8-1.2)	1.9 (1.1-2.8)	3.1 (1.7-4.5)	...
CNS	3.4 (3.1-3.8)	4.2 (2.9-5.4)	6.7 (4.7-8.7)	...
Hemopoietic/lymphatic	3.5 (3.1-3.9)	6.1 (4.6-7.5)	6.4 (4.4-8.3)	...
Solid tumor, non-CNS	3.7 (3.2-4.1)	5.9 (4.5-7.3)	4.1 (2.5-5.7)	4.3 (0.1-8.6)
Metabolic/endocrine				
Diabetes mellitus, insulin dependent	3.7 (3.3-4.1)	2.9 (1.9-3.9)	2.3 (1.1-3.4)	...
Other childhood-onset conditions	2.1 (1.8-2.4)	4.4 (3.2-5.7)	4.9 (3.2-6.6)	5.4 (0.7-10.2)
Neurologic/neurodevelopmental/neuromuscular				
Cerebral palsy	4.1 (3.6-4.5)	7.8 (6.2-9.5)	10.3 (7.9-12.7)	6.5 (1.4-11.7)
Cerebral structural abnormality, congenital	2.4 (2-2.7)	2.8 (1.8-3.8)	4.2 (2.6-5.9)	1.1 (0-3.2)
Developmental delay, generalized/hypoxic ischemic encephalopathy	4.9 (4.4-5.3)	9.1 (7.3-10.8)	13.6 (10.8-16.3)	15.2 (7.7-22.7)
Developmental delay, specific/autism	3.3 (2.9-3.7)	6 (4.5-7.4)	5.4 (3.6-7.2)	6.5 (1.4-11.7)
Epilepsy	7.6 (7.1-8.2)	12 (10-13.9)	16.8 (13.9-19.8)	13 (6-20.1)
Hydrocephalus	4.2 (3.8-4.6)	5.9 (4.5-7.3)	8.3 (6.1-10.5)	5.4 (0.7-10.2)
Muscular dystrophy	1.6 (1.3-1.9)	3.1 (2-4.1)	3.9 (2.4-5.5)	...
Spina bifida/congenital spinal cord abnormality	1 (0.8-1.2)	2.2 (1.3-3.1)	4.1 (2.5-5.7)	2.2 (0-5.2)
Renal/genitourinary				
Chronic renal failure/dialysis dependence	2.2 (1.9-2.5)	6.2 (4.7-7.6)	4.1 (2.5-5.7)	5.4 (0.7-10.2)
Other childhood-onset conditions	1.1 (0.9-1.3)	2.9 (1.9-3.9)	2.5 (1.2-3.7)	1.1 (0-3.2)
Respiratory				
Asthma	6.8 (6.3-7.4)	5 (3.7-6.4)	5.9 (4-7.8)	4.3 (0.1-8.6)
Chronic invasive ventilation	1.1 (0.9-1.3)	2.9 (2-4.1)	5.4 (3.8-7.4)	1 (0-5.4)
Cystic fibrosis	0.7 (0.6-0.9)	1.6 (0.9-2.4)	2 (0.9-3.1)	6.5 (1.4-11.7)
Tracheostomy	3.2 (2.8-3.6)	7 (5.3-8.7)	13.7 (10.8-16.6)	6.5 (1.4-11.7)
Other childhood-onset conditions	1.2 (1-1.5)	3 (2-4)	2.1 (1-3.3)	...
Spine curvature conditions	9.6 (9-10.3)	12.5 (10.4-14.5)	12.3 (9.6-14.9)	8.7 (2.8-14.6)

Abbreviations: CNS, central nervous system; ellipses, no admissions.

<sup>a</sup>Based on 10 481 admissions (96%) of patients 15 years or older from 68 units that supplied secondary diagnoses.

such conditions may preferentially go to PICUs. The reasons are likely multifactorial and similar to the reasons adults use pediatric services in general. Young adults with chronic conditions and their families often have preexisting relationships with pediatric providers and facilities that have expertise and specialized care unique to those conditions.<sup>19,20</sup> They may then be more likely to be admitted for their perioperative and critical care needs to PICUs with which those providers are affiliated. The number of adults in the PICUs may be a proxy for a lack of disease-specific resources and expertise in adult ICUs (eg, perioperative care in congenital heart disease). Other reasons for persistent care of adults in PICUs likely include a lack of planning on the part of pediatric providers,<sup>9,21</sup> a lack of adult-oriented providers with adequate training and experience caring for childhood-onset chronic conditions,<sup>19,20,22,23</sup> a lack of collaboration between pediatric and adult providers, and other factors.<sup>24,25</sup>

The higher mortality of adult PICU patients may be due to progression of certain life-limiting childhood-onset conditions to a terminal stage or, for a smaller segment, adult-onset conditions that impact mortality. These patients may remain under the care of pediatricians for purposes of end-of-life care and despite advanced age. The higher risk-adjusted PICU mortality of adults compared with adolescents and the considerable institutional variability in the proportion of adults have important implications for PICU case mix and benchmarking. Case-mix imbalances between institutions with high or low proportions of adults may unduly impact individual PICUs' predicted mortality and lead to incorrect conclusions regarding outcome performance. A potential solution to case-mix imbalances includes calculating PICU performance for just children. It is unclear if pediatric risk-adjustment systems are valid for adults in PICUs, and adult

**Table 4. Common Acute Conditions by Age Groups**

Condition	% of Admissions (95% CI) <sup>a</sup>			
	Age 15-18 y (n = 8741)	Age 19-20 y (n = 1036)	Age 21-29 y (n = 612)	Age ≥30 y (n = 92)
<b>Cardiac</b>				
Arrhythmia	6 (5.5-6.5)	5.8 (4.4-7.2)	13.4 (10.7-16.1)	21.7 (13.2-30.3)
Congestive heart failure	0.9 (0.7-1.1)	3.3 (2.2-4.4)	3.9 (2.4-5.5)	7.6 (2.1-13.1)
Shock	3.8 (3.4-4.3)	5.2 (3.9-6.6)	5.2 (3.5-7)	7.6 (2.1-13.1)
<b>Gastrointestinal hemorrhage</b>	1.4 (1.2-1.7)	1.4 (0.6-2.1)	2.1 (1-3.3)	2.2 (0-5.2)
<b>Ketoacidosis</b>	9.8 (9.2-10.4)	5.4 (4-6.8)	0.8 (0.1-1.5)	3.3 (0-7)
<b>MODS/SIRS</b>	0.3 (0.2-0.5)	0.8 (0.2-1.3)	1.1 (0.3-2)	2.2 (0-5.2)
<b>Neurologic</b>				
Altered mental status/encephalopathy	7.6 (7-8.1)	7.4 (5.8-9)	4.6 (2.9-6.2)	7.6 (2.1-13.1)
Seizure	4.9 (4.5-5.4)	6.4 (4.9-7.9)	6.4 (4.4-8.3)	4.3 (0.1-8.6)
<b>Poisoning</b>	9.3 (8.7-9.9)	1.8 (1-2.7)	0.8 (0.1-1.5)	...
<b>Renal</b>				
Acute renal failure	2.7 (2.3-3)	4.1 (2.9-5.3)	4.4 (2.8-6)	6.5 (1.4-11.7)
Pyelonephritis/cystitis	1.2 (1-1.4)	1.5 (0.8-2.3)	1.8 (0.7-2.9)	5.4 (0.7-10.2)
<b>Respiratory</b>				
Acute lung injury/ARDS/respiratory failure	11.8 (11.1-12.5)	15.3 (13.1-17.5)	24.8 (21.4-28.3)	19.6 (11.3-27.8)
Pneumonia/pneumonitis	5.4 (5-5.9)	7.6 (6-9.2)	10.1 (7.7-12.5)	14.1 (6.9-21.4)
Status asthmaticus	2.5 (2.2-2.8)	1.4 (0.7-2.2)	1.1 (0.3-2)	1.1 (0-3.2)
<b>Septicemia/bacteremia/viremia</b>	3.5 (3.1-3.9)	7 (5.5-8.6)	5.1 (3.3-6.8)	5.4 (0.7-10.2)

Abbreviations: ARDS, acute respiratory distress syndrome; CSF, cerebral spinal fluid; ellipses, no admissions; MODS, multiple organ dysfunction syndrome; SIRS, systemic inflammatory response syndrome.

<sup>a</sup>Percentages based on 10 481 admissions (96%) of patients 15 years or older from 68 units that supplied secondary diagnoses.

**Table 5. Procedures and Operations by Age Groups**

Condition	% of Admissions (95% CI) <sup>a</sup>			
	Age 15-18 y (n = 5814)	Age 19-20 y (n = 741)	Age 21-29 y (n = 384)	Age ≥30 y (n = 50)
<b>Cardiothoracic</b>				
Assistive device/pacemaker	4.6 (2.9-6.3)	0.2 (0-0.6)	3.4 (1.6-5.2)	10 (1.4-18.6)
Cardiac	3.7 (3.2-4.1)	5.1 (3.5-6.7)	10.2 (7.1-13.2)	32 (18.6-45.4)
Catheterization, therapeutic/diagnostic	2.1 (1.7-2.5)	2.3 (1.2-3.4)	2.1 (0.6-3.5)	2 (0-6)
Thoracic	2.8 (2.3-3.2)	3.8 (2.4-5.2)	3.6 (1.8-5.5)	4 (0-9.6)
<b>Gastrointestinal</b>				
Gastrostomy	0.7 (0.4-0.9)	0.1 (0.2-1.6)	1.6 (0.3-2.8)	2 (0-6)
Other	4.6 (4-5.1)	7 (5.2-8.9)	6.8 (4.2-9.3)	10 (1.4-18.6)
<b>Genitourinary</b>	1.4 (1.1-1.7)	2.6 (1.4-3.7)	1 (0.2-2.1)	8 (0.2-15.8)
<b>Neurologic</b>				
Craniotomy/craniectomy	5.9 (5.3-6.5)	4.6 (3.1-6.1)	6.3 (3.8-8.7)	...
CSF shunt	1.3 (1-1.6)	1.9 (0.9-2.9)	3.6 (1.8-5.5)	2 (0-6)
Other	5.1 (4.5-5.6)	3.8 (2.4-5.2)	4.1 (2.2-6.2)	...
<b>Orthopedic</b>				
Spinal fusion	7.4 (6.8-8.1)	5.8 (4.1-7.5)	1.6 (0.3-2.8)	...
Other	3 (2.6-3.4)	2.7 (1.5-3.9)	1.3 (0.2-2.4)	...
<b>Otolaryngologic</b>				
Tracheotomy	1 (0.7-1.2)	1.3 (0.5-2.2)	2.3 (0.8-3.9)	2 (0-6)
Other	2.1 (1.7-2.4)	2.3 (1.2-3.4)	3.8 (1.6-5.2)	4 (0-9.6)
<b>Vascular</b>	0.7 (0.5-0.9)	0.5 (0-1.1)	1 (0-2.1)	...

Abbreviations: CSF, cerebral spinal fluid; ellipses, no admissions.

<sup>a</sup>Percentages based on 6989 admissions (64%) of patients 15 years or older from 44 units that supplied procedures.

systems<sup>26-28</sup> may not be valid for adults with childhood-onset chronic conditions.

Our study does not answer the questions of if and when all adults should be transitioned or if the quality of care is better in pediatric or adult ICUs. Assuming the survival of persons with childhood-onset chronic conditions will continue to increase, it does suggest that high-

risk adults will continue to be a clinically important patient population, especially for larger PICUs. Pediatric intensive care units, particularly those associated with congenital cardiac programs, should anticipate providing more critical and perioperative care to these patients. At the same time, as efforts to transition these patients to adult care become more widespread and effective, some

of these patients may move from PICUs to adult ICUs. Thus, our study also illustrates the type of patients that adult ICUs will need to prepare for as transition efforts become more successful.

Our study has a number of limitations. First, it is a 1-year cross-sectional study. Therefore, we are unable to comment on trends in the proportion of adult PICU patients. The proportion of adults in our sample may have decreased with advancing age for various unstudied reasons—transition to adult care, death, and relocation, for example. Second, the PICUs included in our database may not be representative of all PICUs in the United States. Because institutional data were only provided in the aggregate, we could not stratify by types of PICUs. Also, a limited number of data variables were not submitted by every PICU, although the vast majority of information was captured completely. Third, any postulations on the relative appropriateness of admitting adults to PICUs based on these findings should be made in light of the fact that we do not know what adult critical care options, if any, these patients may have had. Similarly, we do not have data on adults with childhood-onset chronic conditions who use adult ICUs. Therefore, we do not know how the characteristics of our adult PICU patients compare with those of the overall population of adults with childhood-onset conditions needing ICU care. Lastly, our estimate of the increased risk of PICU mortality associated with age 21 years or older was adjusted using a pediatric severity-of-illness scoring system. Neither Paediatric Index of Mortality 2 score nor Pediatric Risk of Mortality score<sup>29</sup>—the 2 most widely used pediatric severity-of-illness systems—have been validated for adults. Nevertheless, PICUs are using these systems to risk adjust outcomes for their adult patients.

## CONCLUSIONS

The combination of increasing survival of young adults with chronic conditions and possibly underdeveloped transitioning efforts has resulted in a small, but clinically important, cohort of adults with critical illness who receive care in PICUs. Characterizing the extent and patterns of pediatric critical care use by adults is helpful for planning both the provision of future clinical services and further research to evaluate the success of transition efforts. Additional research is needed to examine the trends of adult PICU admissions, the reasons for institutional variation, how to effectively transition suitable adult populations from pediatric to adult critical care services, their costs of care, and if the quality of care is different for these adults when cared for in adult ICUs.

It is unclear if PICUs or adult ICUs are the best place to meet the critical care needs of adults with childhood-onset chronic conditions, and certain conditions may be better cared for in particular units. Pediatric facilities, including PICUs, often do not have the specific resources and expertise needed to optimize the care of adult-acquired conditions. On the other hand, PICUs may be more capable than adult ICUs of providing the specialized care needed to address their underlying childhood-onset chronic conditions. Regardless, as the number of

adults with childhood-onset chronic conditions grows, PICUs will need to prepare for increasingly older patients and adult ICUs will increasingly need to care for these adults. Both types of units will need to make greater efforts to collaborate on research and training of providers to meet their unique care needs and on guidelines that help direct specific types of adult patients to the unit type where those needs can best be met.

**Accepted for Publication:** October 30, 2012.

**Published Online:** March 18, 2013. doi:10.1001/jamapediatrics.2013.1316

**Author Affiliations:** Division of Pediatric Critical Care Medicine, Department of Pediatrics, Columbia University College of Physicians and Surgeons, New York, New York (Dr Edwards); Department of Physical Medicine and Rehabilitation, University of Pittsburgh, Pittsburgh, Pennsylvania (Dr Houtrow); Division of General Internal Medicine and Public Health and Center for Health Services Research, Vanderbilt University School of Medicine (Dr Vasilevskis), and Geriatric Research Education and Clinical Center Service and Clinical Research Training Center of Excellence, Department of Veterans Affairs Medical Center, Tennessee Valley Healthcare System (Dr Vasilevskis), Nashville; and Divisions of Pulmonary and Critical Care (Dr Dudley) and General Internal Medicine (Dr Okumura), Department of Medicine, Philip R. Lee Institute for Health Policy Studies (Drs Dudley and Okumura), and Division of General Pediatrics, Department of Pediatrics (Dr Okumura), University of California, San Francisco.

**Correspondence:** Jeffrey D. Edwards, MD, MA, MAS, Division of Pediatric Critical Care Medicine, Columbia University College of Physicians and Surgeons, 3959 Broadway, CHN 10-24, New York, NY 10032 (jde2134@columbia.edu).

**Author Contributions:** *Study concept and design:* Edwards and Okumura. *Acquisition of data:* Edwards. *Analysis and interpretation of data:* All authors. *Drafting of the manuscript:* Edwards. *Critical revision of the manuscript for important intellectual content:* All authors. *Statistical analysis:* Edwards. *Obtained funding:* Edwards. *Administrative, technical, and material support:* Okumura. *Study supervision:* Houtrow, Dudley, and Okumura.

**Conflict of Interest Disclosures:** None reported.

**Funding/Support:** Dr Edwards is supported by National Institutes of Health (NIH) Pediatric Critical Care Scientist Development Program grant K12 HD 047349. Dr Houtrow is supported by NIH grant K12 H001097. Dr Vasilevskis is supported by NIH grant K23 AG040157 and receives support from the Geriatric Research Education and Clinical Centers and the Clinical Research Training Center of Excellence, Department of Veterans Affairs Medical Center, Tennessee Valley Healthcare System, Nashville. Dr Okumura is supported by Agency for Healthcare Research and Quality grant K08 HS017716.

**Additional Contributions:** We thank the Virtual Pediatric Intensive Care Unit Systems for providing the data for this study. We also thank the Pediatric Critical Care Scientist Development Program for the grant support that permitted this work. Finally, we thank Harsha Reddy, MD, for review of the manuscript.

## REFERENCES

1. American Academy of Pediatrics; American Academy of Family Physicians; American College of Physicians-American Society of Internal Medicine. A consensus statement on health care transitions for young adults with special health care needs. *Pediatrics*. 2002;110(6, pt 2):1304-1306.
2. Rosen DS, Blum RW, Britto M, Sawyer SM, Siegel DM; Society for Adolescent Medicine. Transition to adult health care for adolescents and young adults with chronic conditions: position paper of the Society for Adolescent Medicine. *J Adolesc Health*. 2003;33(4):309-311.
3. Achieving and measuring success: a national agenda for children with special health care needs. US Dept of Health and Human Services, Maternal and Child Health Bureau website. <http://www.mchb.hrsa.gov/programs/specialneeds/achievinguccess.html>. Accessed February 10, 2012.
4. Cooley WC, Sagerman PJ; American Academy of Pediatrics; American Academy of Family Physicians; American College of Physicians; Transitions Clinical Report Authoring Group. Supporting the health care transition from adolescence to adulthood in the medical home. *Pediatrics*. 2011;128(1):182-200.
5. Burdo-Hartman WA, Patel DR. Medical home and transition planning for children and youth with special health care needs. *Pediatr Clin North Am*. 2008;55(6):1287-1297, vii-viii.
6. Goodman DM, Mendez E, Throop C, Ogata ES. Adult survivors of pediatric illness: the impact on pediatric hospitals. *Pediatrics*. 2002;110(3):583-589.
7. Okumura MJ, Campbell AD, Nasr SZ, Davis MM. Inpatient health care use among adult survivors of chronic childhood illnesses in the United States. *Arch Pediatr Adolesc Med*. 2006;160(10):1054-1060.
8. Tan J, Cox R, Shannon P, Payne D. Adolescent and young adult health in a children's hospital: everybody's business. *J Paediatr Child Health*. 2009;45(12):715-719.
9. Lam PY, Fitzgerald BB, Sawyer SM. Young adults in children's hospitals: why are they there? *Med J Aust*. 2005;182(8):381-384.
10. Goodman DM, Hall M, Levin A, et al. Adults with chronic health conditions originating in childhood: inpatient experience in children's hospitals. *Pediatrics*. 2011;128(1):5-13.
11. McDonnell WM, Kocolas I, Roosevelt GE, Yetman AT. Pediatric emergency department use by adults with chronic pediatric disorders. *Arch Pediatr Adolesc Med*. 2010;164(6):572-576.
12. Baker MD, Schwartz GR, Ludwig S. The adult patient in the pediatric emergency department. *Ann Emerg Med*. 1993;22(7):1136-1139.
13. Slater A, Shann F, Pearson G; Paediatric Index of Mortality (PIM) Study Group. PIM2: a revised version of the Paediatric Index of Mortality. *Intensive Care Med*. 2003;29(2):278-285.
14. Feudtner C, Christakis DA, Connell FA. Pediatric deaths attributable to complex chronic conditions: a population-based study of Washington State, 1980-1997. *Pediatrics*. 2000;106(1, pt 2):205-209.
15. Edwards JD, Houtrow AJ, Vasilevskis EE, et al. Chronic conditions among children admitted to U.S. pediatric intensive care units: their prevalence and impact on risk for mortality and prolonged length of stay. *Crit Care Med*. 2012;40(7):2196-2203.
16. Fiser DH. Assessing the outcome of pediatric intensive care. *J Pediatr*. 1992;121(1):68-74.
17. Fact sheets. National Association of Children's Hospitals and Related Institutions/ National Association of Children's Hospitals website. [http://www.childrenshospitals.net/AM/Template.cfm?Section=Facts\\_and\\_Trends](http://www.childrenshospitals.net/AM/Template.cfm?Section=Facts_and_Trends). Accessed February 10, 2012.
18. Rosenberg DI, Moss MM; American Academy of Pediatrics Section on Critical Care; American Academy of Pediatrics Committee on Hospital Care. Guidelines and levels of care for pediatric intensive care units. *Pediatrics*. 2004;114(4):1114-1125.
19. Reiss J, Gibson RW. At what cost? *Arch Pediatr Adolesc Med*. 2006;160(10):1079-1080.
20. Reiss JG, Gibson RW, Walker LR. Health care transition: youth, family, and provider perspectives. *Pediatrics*. 2005;115(1):112-120.
21. Lotstein DS, Ghandour R, Cash A, McGuire E, Strickland B, Newacheck P. Planning for health care transitions: results from the 2005-2006 National Survey of Children With Special Health Care Needs. *Pediatrics*. 2009;123(1):e145-e152.
22. Patel MS, O'Hare K. Residency training in transition of youth with childhood-onset chronic disease. *Pediatrics*. 2010;126(suppl 3):S190-S193.
23. Brennan LJ, Rolfe PM. Transition from pediatric to adult health services: the perioperative care perspective. *Paediatr Anaesth*. 2011;21(6):630-635.
24. Peter NG, Forke CM, Ginsburg KR, Schwarz DF. Transition from pediatric to adult care: internists' perspectives. *Pediatrics*. 2009;123(2):417-423.
25. Okumura MJ, Kerr EA, Cabana MD, Davis MM, Demonner S, Heisler M. Physician views on barriers to primary care for young adults with childhood-onset chronic disease. *Pediatrics*. 2010;125(4):e748-e754.
26. Zimmerman JE, Kramer AA, McNair DS, Malila FM. Acute Physiology and Chronic Health Evaluation (APACHE) IV: hospital mortality assessment for today's critically ill patients. *Crit Care Med*. 2006;34(5):1297-1310.
27. Higgins TL, Kramer AA, Nathanson BH, Copes W, Stark M, Teres D. Prospective validation of the intensive care unit admission Mortality Probability Model (MPMO-III). *Crit Care Med*. 2009;37(5):1619-1623.
28. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-383.
29. Pollack MM, Patel KM, Ruttimann UE. PRISM III: an updated Pediatric Risk of Mortality score. *Crit Care Med*. 1996;24(5):743-752.